

Letter to the Editor

Comment on “Web-Based Measure of Life Events Using Computerized Life Events and Assessment Record (CLEAR): Preliminary Cross-Sectional Study of Reliability, Validity, and Association With Depression”: Validity and Methodological Issues

Jamal Rahmani¹, MSc, PhD; Roya Karimi², MSc, PhD; Farideh Mohtasham³, MSc, PhD; Siamak Sabour^{2,4}, MD, PhD

¹National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Department of Clinical Epidemiology, School of Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³Department of Clinical Epidemiology, School of Health and Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Safety Promotions and Injury Prevention Research Centre, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Corresponding Author:

Siamak Sabour, MD, PhD

Department of Clinical Epidemiology

School of Health and Safety

Shahid Beheshti University of Medical Sciences

Chamran Highway

Tehran

Iran

Phone: 98 21 22421814

Email: s.sabour@sbmu.ac.ir

Related Articles:

Comment on: <https://mental.jmir.org/2019/1/e10675/>

Comment in: <https://mental.jmir.org/2020/5/e15434/>

(*JMIR Ment Health* 2020;7(5):e14505) doi: [10.2196/14505](https://doi.org/10.2196/14505)

KEYWORDS

validity; methodological issues; diagnostic test

We were interested in the article titled, “Web-Based Measure of Life Events Using Computerized Life Events and Assessment Record (CLEAR): Preliminary Cross-Sectional Study of Reliability, Validity, and Association With Depression” published in *JMIR Mental Health* [1].

One of the aims of the abovementioned study was to assess the validity of Computerized Life Events and Assessment Record (CLEAR), considering the Life Events and Difficulties Schedule (LEDS) and the List of Threatening Experiences Questionnaire (LTE-Q) as gold standards among 328 participants (126 students; 202 matched midlife sample: 127 unaffected controls, 75 recurrent depression cases). The authors concluded that CLEAR has acceptable validity and great potential for effective use in research and clinical practice. However, there are some methodological issues in this conclusion that are mentioned below.

First, there are some measures that can be applied to the assessment of the validity of a test including sensitivity, specificity, positive predictive value, negative predictive value,

positive likelihood ratio (LR+; ranging from 1 to infinity; the higher the LR+, the more accurate is the test), negative likelihood ratio (LR-; ranging from 0 to 1; the lower the LR-, the more accurate is the test), and odds ratio (ratio of true to false results) [2-5]. According to the results, sensitivity of CLEAR was 59.1% and 43.1% compared to LEDS and LTE-Q, respectively, as gold standards. Likewise, specificity of CLEAR was 65.4% and 78.6%, respectively, compared to the abovementioned gold standards.

It is good to know that sensitivity is an important measure in public health aspects instead of clinical fields. Likewise, the positive predictive value and negative predictive value are among measures that are more appropriate for advice about the validity of a diagnostic test for clinical purposes [3-5]. Therefore, we suggest applying predictive values, likelihood ratios, odds ratio, and diagnostic accuracy to decide the validity of CLEAR. Moreover, according to the data of study, LR+, LR-, odds ratio, and diagnostic accuracy of CLEAR will be 1.6, 0.6, 2.6, and 62%, respectively, compared to LEDS (Tables 1 and 2) and 1.9, 0.7, 2.6, and 60%, respectively compared to

LTE-Q (Tables 3 and 4). Therefore, there is a high level of insufficient evidence to conclude that the validity of the CLEAR uncertainty for decisions based on these values, and there is test is acceptable.

Table 1. Two by two table of Computerized Life Events and Assessment Record compared to Life Events and Difficulties Schedule as the gold standard.

CLEAR ^a	LEDS ^b (gold standard)		
	Positive	Negative	Total
Positive	59	35	94
Negative	41	65	106
Total	100	100	200

^aCLEAR: Computerized Life Events and Assessment Record.

^bLEDS: Life Events and Difficulties Schedule.

Table 2. Assessing the validity of Computerized Life Events and Assessment Record compared to Life Events and Difficulties Schedule as the gold standard.

Parameter	Estimate
Sensitivity (%)	59
Specificity (%)	65
Positive predictive value (%)	63
Negative predictive value (%)	61
Diagnostic accuracy (%)	62
Likelihood ratio of a positive test	1.6
Likelihood ratio of a negative test	0.6
Diagnostic odds	2.6

Table 3. Two by two table of Computerized Life Events and Assessment Record compared to List of Threatening Experiences Questionnaire as the gold standard.

CLEAR ^a	LTE-Q ^b (gold standard)		
	Positive	Negative	Total
Positive	43	22	65
Negative	57	78	135
Total	100	100	200

^aCLEAR: Computerized Life Events and Assessment Record.

^bLTE-Q: List of Threatening Experiences Questionnaire.

Table 4. Assessing the validity of Computerized Life Events and Assessment Record compared to List of Threatening Experiences Questionnaire as the gold standard.

Parameter	Estimate
Sensitivity (%)	43
Specificity (%)	78
Positive predictive value (%)	66
Negative predictive value (%)	58
Diagnostic accuracy (%)	60
Likelihood ratio of a positive test	1.9
Likelihood ratio of a negative test	0.7
Diagnostic odds	2.6

Conflicts of Interest

None declared.

References

1. Bifulco A, Spence R, Nunn S, Kagan L, Bailey-Rodriguez D, Hosang GM, et al. Web-Based Measure of Life Events Using Computerized Life Events and Assessment Record (CLEAR): Preliminary Cross-Sectional Study of Reliability, Validity, and Association With Depression. *JMIR Ment Health* 2019 Jan 08;6(1):e10675 [FREE Full text] [doi: [10.2196/10675](https://doi.org/10.2196/10675)] [Medline: [30622088](https://pubmed.ncbi.nlm.nih.gov/30622088/)]
2. Szklo M, Nieto J. *Epidemiology: beyond the basics*. 3rd ed. Manhattan, NY: Jones and Bartlett Learning; 2014.
3. Sabour S, Ghassemi F. The validity and reliability of a signal impact assessment tool: statistical issue to avoid misinterpretation. *Pharmacoepidemiol Drug Saf* 2016 Oct;25(10):1215-1216. [doi: [10.1002/pds.4061](https://doi.org/10.1002/pds.4061)] [Medline: [27696610](https://pubmed.ncbi.nlm.nih.gov/27696610/)]
4. Sabour S. Validity and reliability of the new Canadian Nutrition Screening Tool in the 'real-world' hospital setting: Methodological issues. *Eur J Clin Nutr* 2015 Jul;69(7):864. [doi: [10.1038/ejcn.2015.69](https://doi.org/10.1038/ejcn.2015.69)] [Medline: [25920428](https://pubmed.ncbi.nlm.nih.gov/25920428/)]
5. Sabour S. Validity and reliability of the 13C-methionine breath test for the detection of moderate hyperhomocysteinemia in Mexican adults; statistical issues in validity and reliability analysis. *Clin Chem Lab Med* 2014 Dec;52(12):e295-e296. [doi: [10.1515/cclm-2014-0453](https://doi.org/10.1515/cclm-2014-0453)] [Medline: [24933629](https://pubmed.ncbi.nlm.nih.gov/24933629/)]

Abbreviations

CLEAR: Computerized Life Events and Assessment Record

LEDS: Life Events and Difficulties Schedule

LR-: negative likelihood ratio

LR+: positive likelihood ratio

LTE-Q: List of Threatening Experiences Questionnaire

Edited by J Torous; submitted 26.04.19; peer-reviewed by A Bifulco; accepted 10.07.19; published 21.05.20

Please cite as:

Rahmani J, Karimi R, Mohtasham F, Sabour S

Comment on "Web-Based Measure of Life Events Using Computerized Life Events and Assessment Record (CLEAR): Preliminary Cross-Sectional Study of Reliability, Validity, and Association With Depression": Validity and Methodological Issues
JMIR Ment Health 2020;7(5):e14505

URL: <https://mental.jmir.org/2020/5/e14505>

doi: [10.2196/14505](https://doi.org/10.2196/14505)

PMID:

©Jamal Rahmani, Roya Karimi, Farideh Mohtasham, Siamak Sabour. Originally published in *JMIR Mental Health* (<http://mental.jmir.org>), 21.05.2020. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR Mental Health*, is properly cited. The complete bibliographic information, a link to the original publication on <http://mental.jmir.org/>, as well as this copyright and license information must be included.